



Chinese Weekly Influenza Surveillance Report

June 29 to July 5, 2026 (Week 27)

(All data are preliminary and may change as more reports are received)

Summary

- The positive rate of influenza virus testing was increased in southern provinces, while decreased in northern provinces. Influenza B predominated. There were 7 ILI outbreaks reported in this week.
- Among influenza viruses antigenically characterized by CNIC since March 30, 2026, 18(100.0%) influenza A(H1N1)pdm09 viruses were characterized as A/Missouri/11/2025-like; 139(76.4%) influenza A(H3N2) viruses were characterized as A/Darwin/1454/2025(egg)-like, 161(88.5%) influenza A(H3N2) viruses were characterized as A/Darwin/1415/2025(cell)-like; 349(94.1%) influenza B/Victoria viruses were characterized as B/Tokyo/EIS13-175/2025-like.
- Among the influenza viruses tested by CNIC for antiviral resistance analysis since March 30, 2026, all A(H1N1)pdm09, A(H3N2) and B viruses were sensitive to neuraminidase inhibitors; all A(H1N1)pdm09, A(H3N2) and B viruses were sensitive to influenza polymerase inhibitors.

Surveillance of outpatient or emergency visits for Influenza-like Illness (ILI)

During week 27, the percentage of outpatient or emergency visits for ILI (ILI%) at national sentinel hospitals in southern provinces was 5.6%, higher than the last week (5.4%), higher than the same week of 2023 and 2025(4.8% and 4.2%), the same as the same week of 2024(5.6%). (Figure 1)




Figure 1. Percentage of Visits for ILI at Sentinel Hospitals in Southern Provinces

Note: Analysis in this part was based on data from sentinel hospitals belong to national influenza surveillance network.

During week 27, ILI% at national sentinel hospitals in northern provinces was 3.0%, the same as the last week (3.0%), lower than the same week of 2024(3.7%), higher than the same week of 2025(2.8%). (Figure 2)




Figure 2. Percentage of Visits for ILI at Sentinel Hospitals in Northern Provinces

Note: Analysis in this part was based on data from sentinel hospitals belong to national influenza surveillance network.



Virologic Surveillance

During week 27 of 2026, influenza network laboratories tested 13676 specimens, there were 1639 positive detections for influenza. The number and proportion of influenza types and subtypes detected in southern and northern provinces were shown in Table 1.

Table 1. Laboratory Detections of ILI Specimens (Week 27, 2026)

	Week 27		
	Southern provinces	Northern provinces	Total
No. of specimens tested	10988	2688	13676
No. of positive specimens (%)	1579(14.4%)	60(2.2%)	1639(12.0%)
Influenza A	469(29.7%)	13(21.7%)	482(29.4%)
A(H1N1)pdm09	19(4.1%)	2(15.4%)	21(4.4%)
A(H3N2)	450(95.9%)	11(84.6%)	461(95.6%)
A (subtype not determined)	0	0	0
Influenza B	1110(70.3%)	47(78.3%)	1157(70.6%)
B (lineage not determined)	0	0	0
Victoria	1110(100.0%)	47(100.0%)	1157(100.0%)
Yamagata	0	0	0




Figure 3. Influenza Positive Tests Reported by Southern Network Laboratories

Note: Analysis in this part was based on the test results of network laboratories. If it were not consistent with the results of CNIC confirmation, the results of CNIC confirmation were used.




Figure 4. Influenza Positive Tests Reported by Northern Network Laboratories

Note: Analysis in this part was based on the result of network laboratories. If it were not consistent with the results of CNIC confirmation, the results of CNIC confirmation were used.

Antigenic Characterization

Since March 30, 2026, 18(100.0%) influenza A(H1N1)pdm09 viruses were characterized as A/Missouri/11/2025-like; 139(76.4%) influenza A(H3N2) viruses were characterized as A/Darwin/1454/2025(egg)-like, 161(88.5%) influenza A(H3N2)viruses were characterized as A/Darwin/1415/2025(cell)-like; 349(94.1%) influenza B/Victoria viruses were characterized as B/Tokyo/EIS13-175/2025-like.



Antiviral Resistance

Since March 30, 2026, among the influenza viruses tested by CNIC for antiviral resistance, all A(H1N1)pdm09, A(H3N2) and B viruses were sensitive to neuraminidase inhibitors; all A(H1N1)pdm09, A(H3N2) and B viruses were sensitive to influenza polymerase inhibitors.

Outbreak Surveillance

During week 27, there were 7 ILI outbreaks reported nationwide. Among them, 3 were B/Victoria, 2 were A(H3N2), 1 was mixed, 1 was negative for flu.